

April 14, 1999

UCB-PTH-99/17

LBNL-43119

hep-th/9904105

General Virasoro Construction on Orbifold Affine Algebra

J. EVSLIN, M. B. HALPERN, and J. E. WANG*

*Department of Physics, University of California
Berkeley, California 94720*

and

*Theoretical Physics Group
Ernest Orlando Lawrence Berkeley National Laboratory
University of California, Berkeley, California 94720*

Abstract

We obtain the orbifold Virasoro master equation (OVME) at integer order λ , which summarizes the general Virasoro construction on orbifold affine algebra. The OVME includes the Virasoro master equation when $\lambda = 1$ and contains large classes of stress tensors of twisted sectors of conventional orbifolds at higher λ . The generic construction is like a twisted sector of an orbifold (with non-zero ground state conformal weight) but new constructions are obtained for which we have so far found no conventional orbifold interpretation.

*E-Mail: hllywd2@physics.berkeley.edu

1 Introduction

The Virasoro master equation (VME) summarizes the general Virasoro construction^{1,2} on affine Lie algebra,

$$T(z) = L^{ab} : J_a(z) J_b(z) : , \quad a, b = 1, \dots, \dim g . \quad (1.1)$$

See Ref. 3 for a review of the VME and irrational conformal field theory. Subsequent developments in this subject include semiclassical examples of conformal blocks⁴ in irrational conformal field theory, unification⁵⁻⁸ with the general non-linear sigma model and the construction of the stress tensors of twisted sectors of cyclic permutation orbifolds⁹ from copies of the general Virasoro construction (1.1).

Our starting point in this paper is the orbifold affine algebra⁹ at integer order λ , which was recently found in the twisted sectors of cyclic permutation orbifolds and includes affine Lie algebra when $\lambda = 1$. Following the suggestion of Ref. 9, we consider here the general Virasoro construction on orbifold affine algebra

$$\hat{T}(z) = \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)}(z) \hat{J}_b^{(-r)}(z) : , \quad a, b = 1, \dots, \dim g, \quad \lambda \in \mathbb{Z}^+ \quad (1.2)$$

where $\hat{J}_a^{(r)}(z)$, $r = 0, \dots, \lambda - 1$ are the orbifold currents. When $\lambda = 1$, the orbifold currents $\hat{J}_a^{(0)}(z) = J_a(z)$ satisfy affine Lie algebra and $\hat{T}(z)$ reduces to $T(z)$ in (1.1).

The stress tensor $\hat{T}(z)$ is conformal when $\mathcal{L}_r^{ab}$ satisfies the *orbifold Virasoro master equation* (OVME) at order λ , and the OVME contains the conformal field theories of the VME when $\lambda = 1$. At higher λ , the OVME contains large classes of stress tensors of twisted sectors of conventional permutation orbifolds, and many other constructions. The generic construction of the OVME is apparently a twisted sector of an orbifold or a Ramond (spin) sector because the ground state conformal weights $\hat{\Delta}_0$ of the constructions

$$L(m \geq 0)|0\rangle = \delta_{m,0} \hat{\Delta}_0 |0\rangle, \quad \hat{\Delta}_0 \neq 0 \quad (1.3)$$

are generically non-zero. New unitary solutions of the OVME are obtained however, with irrational central charge and/or irrational conformal weights, for which we have so far found no conventional orbifold interpretation.

2 Background

2.1 Orbifold affine algebra

We consider the *orbifold affine algebra*⁹ g_λ

$$[\hat{J}_a^{(r)}(m + \frac{r}{\lambda}), \hat{J}_b^{(s)}(n + \frac{s}{\lambda})] = if_{ab}{}^c \hat{J}_c^{(r+s)}(m+n + \frac{r+s}{\lambda}) + \hat{G}_{ab}(m + \frac{r}{\lambda}) \delta_{m+n+\frac{r+s}{\lambda}, 0} \quad (2.1a)$$

$$m, n \in \mathbb{Z}, \quad a, b = 1, \dots, \dim g, \quad r, s = 0, \dots, \lambda - 1 \quad (2.1b)$$

$$\hat{J}_a^{(r)}(m + \frac{r}{\lambda})|0\rangle = 0 \quad \text{when} \quad m + \frac{r}{\lambda} \geq 0 \quad (2.1c)$$

where the order λ of g_λ is any positive integer and $g_{\lambda=1}$ is affine Lie algebra¹⁰⁻¹². The relations in (2.1) are understood with the periodicity condition

$$\hat{J}_a^{(r \pm \lambda)}(m + \frac{r \pm \lambda}{\lambda}) = \hat{J}_a^{(r)}(m \pm 1 + \frac{r}{\lambda}) \quad (2.2)$$

and the state $|0\rangle$ is called the ground state of g_λ . The quantities $f_{ab}{}^c$ are the structure constants of any semisimple Lie algebra $g = \oplus_I g_I$, and the metric $\hat{G}_{ab}$ is

$$\hat{G}_{ab} = \oplus_I \hat{k}_I \eta_{ab}^I. \quad (2.3)$$

Here, η_{ab}^I is the Killing metric of g_I and $\hat{k}_I = \lambda k_I$ is the level of the associated orbifold affine subalgebra. For applications we restrict ourselves to simple compact g ,

$$\hat{G}_{ab} = \hat{k} \eta_{ab}, \quad \hat{k} = \lambda k, \quad \hat{x} = \frac{2\hat{k}}{\psi^2} = \lambda x, \quad x = \frac{2k}{\psi^2} \quad (2.4)$$

where ψ is the highest root of g and unitarity⁹ requires that the invariant level x be a positive integer.

We will also need the operator product form⁹ of g_λ

$$\hat{J}_a^{(r)}(z) \hat{J}_b^{(s)}(w) = \frac{\hat{G}_{ab} \delta_{r+s, 0 \bmod \lambda}}{(z-w)^2} + \frac{if_{ab}{}^c \hat{J}_c^{(r+s)}(w)}{(z-w)} + O((z-w)^0) \quad (2.5a)$$

$$\hat{J}_a^{(r)}(z) = \sum_{m \in \mathbb{Z}} \hat{J}_a^{(r)}(m + \frac{r}{\lambda}) z^{-1-m-r/\lambda} \quad (2.5b)$$

where $\hat{J}_a^{(r)}(z)$ are the local orbifold currents.

2.2 Subalgebras of g_λ

The algebra g_λ contains a set of regularly embedded subalgebras $g_\eta \subset g_\lambda$,

$$\hat{J}_a^{(r)}(m + \frac{r}{\eta}) = \hat{J}_a^{(\lambda r/\eta)}(m + \frac{\lambda r/\eta}{\lambda}), \quad r = 0, \dots, \eta - 1, \quad \eta \in \mathbb{Z}^+, \quad \frac{\lambda}{\eta} \in \mathbb{Z}^+ \quad (2.6a)$$

$$[\hat{J}_a^{(r)}(m + \frac{r}{\eta}), \hat{J}_b^{(s)}(n + \frac{s}{\eta})] = i f_{ab}^c \hat{J}_c^{(r+s)}(m+n + \frac{r+s}{\eta}) + \hat{G}_{ab}(m + \frac{r}{\eta}) \delta_{m+n+\frac{r+s}{\eta}, 0} \quad (2.6b)$$

which is isomorphic to the set of order η orbifold affine algebras, taken at levels $\{\hat{k}_I = \lambda k_I\}$.

The special case $\eta = 1$ is known as the integral affine subalgebra^{13,14,9}, whose generators

$$\hat{J}_a^{(0)}(m) = \hat{J}_a^{(0)}(m) \quad (2.7)$$

form an affine Lie algebra at levels $\{\hat{k}_I\}$. Related subalgebras include $h_\eta \subset g_\lambda$, $\lambda/\eta \in \mathbb{Z}^+$ where $h \subset g$ is any Lie subalgebra of g .

There are many other subalgebras of g_λ , for example the algebra generated by

$$\hat{J}_a^{(\mu n \bmod \lambda)}(\lfloor \frac{\mu n}{\lambda} \rfloor + \frac{\mu n \bmod \lambda}{\lambda}), \quad n \in \mathbb{Z}, \quad \mu = 0, 1, 2, \dots \quad (2.8)$$

where $\lfloor x \rfloor$ is the integer less than or equal to x . When $\mu = \frac{\lambda}{\eta} \in \mathbb{Z}^+$, these are the generators

$$\hat{J}_a^{(r)}(m + \frac{r}{\eta}) = \hat{J}_a^{(\mu r)}(m + \frac{\mu r}{\lambda}), \quad r = 0, \dots, \eta - 1 \quad (2.9)$$

of the g_η subalgebras above.

2.3 Normal ordering

In Ref. 9, the (mother theory) normal-ordering convention

$$\begin{aligned} : \hat{A}^{(r)}(z) \hat{B}^{(s)}(z) :_M &\equiv \sum_{m=-\infty}^{\infty} z^{-\Delta(A)-\Delta(B)-m-\frac{r+s}{\lambda}} \left[\sum_{p \leq -\Delta(A)} \hat{A}^{(r)}(p + \frac{r}{\lambda}) \hat{B}^{(s)}(m - p + \frac{s}{\lambda}) \right. \\ &\quad \left. + \sum_{p > -\Delta(A)} \hat{B}^{(s)}(m - p + \frac{s}{\lambda}) \hat{A}^{(r)}(p + \frac{r}{\lambda}) \right] \end{aligned} \quad (2.10)$$

was used for the product of any two integer-moded orbifold principal primary fields, including the orbifold currents with (mother theory) conformal weights $\Delta(A) = \Delta(B) = 1$. We will use instead the OPE normal-ordering convention

$$: \hat{A}^{(r)}(z) \hat{B}^{(s)}(z) : \equiv \oint_z \frac{(dx)}{x - z} \hat{A}^{(r)}(x) \hat{B}^{(s)}(z), \quad (dx) \equiv \frac{dx}{2\pi i} \quad (2.11)$$

where the x contour does not encircle the origin. In the case of the orbifold currents, the two conventions are related by

$$: \hat{J}_a^{(r)}(z) \hat{J}_b^{(s)}(z) : \equiv : \hat{J}_a^{(r)}(z) \hat{J}_b^{(s)}(z) :_M - \frac{ir f_{ab}{}^c \hat{J}_c^{(r+s)}(z)}{\lambda z} + \frac{r(\lambda - r)}{2\lambda^2 z^2} \hat{G}_{ab} \delta_{r+s,0 \bmod \lambda} \quad (2.12)$$

and we note that the modes of the OPE normal-ordered bilinears satisfy

$$: \hat{J}_a^{(r)}(z) \hat{J}_b^{(-r)}(z) : = \sum_{m \in \mathbb{Z}} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : (m) z^{-m-2} \quad (2.13a)$$

$$: \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : (m \geq 0) |0\rangle = \frac{r(\lambda - r)}{2\lambda^2} \hat{G}_{ab} \delta_{m,0} |0\rangle, \quad r = 0, \dots, \lambda - 1 \quad (2.13b)$$

because $: \hat{J}_a^{(r)} \hat{J}_b^{(-r)} :_M$ ($m \geq 0$) annihilates the ground state.

The current bilinears also have a symmetry (see App. A)

$$: \hat{J}_a^{(r)}(z) \hat{J}_b^{(s)}(z) : = : \hat{J}_a^{(s)}(z) \hat{J}_b^{(r)}(z) : \quad (2.14)$$

which will play an important role in the general construction below. (The symmetry holds for $: JJ :_M$ as well.)

3 General Virasoro Construction

3.1 Summary of the computation

Following the suggestion of Ref. 9, we consider the candidate stress tensors

$$\hat{T}(z) = \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)}(z) \hat{J}_b^{(-r)}(z) : = \sum_{m \in \mathbb{Z}} L(m) z^{-m-2} \quad (3.1a)$$

$$\mathcal{L}_r^{ab} = \mathcal{L}_r^{ba}, \quad r = 0, \dots, \lambda - 1 \quad (3.1b)$$

where $\{\mathcal{L}_r^{ab}\}$ is a set of λ “inverse inertia tensors”. Then, following Ref. 1, we require that $\hat{T}$ satisfies the Virasoro algebra

$$\hat{T}(z) \hat{T}(w) = \frac{\hat{c}/2}{(z-w)^4} + \frac{2\hat{T}(w)}{(z-w)^2} + \frac{\partial_w \hat{T}(w)}{(z-w)} + O((z-w)^0) \quad (3.2)$$

This computation (see App. A) results in the restriction on $\mathcal{L}_r^{ab}$

$$\mathcal{F}_r^{ab} = \mathcal{F}_r^{ac} \hat{G}_{cd} \mathcal{F}_r^{db} - \frac{1}{2} \sum_{s=0}^{\lambda-1} \mathcal{F}_s^{cd} [\mathcal{F}_{r+s}^{ef} f_{ce}^a f_{df}^b + f_{ce}^f f_{df}^{(a} \mathcal{F}_r^{b)e}] \quad (3.3a)$$

$$\mathcal{F}_r^{ab} \equiv \mathcal{L}_r^{ab} + \mathcal{L}_{\lambda-r}^{ab}, \quad r = 0, \dots, \lambda - 1 \quad (3.3b)$$

(where we have defined $\mathcal{L}_{r\pm\lambda}^{ab} \equiv \mathcal{L}_r^{ab}$) and the central charge,

$$\hat{c} = 2\hat{G}_{ab} \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab}. \quad (3.4)$$

The form of the restriction (3.3), in terms of the combination $\mathcal{F}_r^{ab}$, is a consequence of the symmetry (2.14). Any solution of the system (3.3) gives a conformal stress tensor $\hat{T}$, but this system is problematic because it has more unknowns than equations.

3.2 Equivalent Solutions

The system (3.3) determines only the combination $\mathcal{F}_r^{ab}$ in (3.3b). This means that (3.3) has a “gauge invariance” under the gauge transformation

$$\mathcal{L}_r^{ab} \rightarrow (\mathcal{L}'_r)^{ab} = \mathcal{L}_r^{ab} + \Lambda_r^{ab}, \quad \mathcal{L}_{\lambda-r}^{ab} \rightarrow (\mathcal{L}'_{\lambda-r})^{ab} = \mathcal{L}_{\lambda-r}^{ab} - \Lambda_r^{ab} \quad (3.5a)$$

$$\mathcal{F}_r^{ab} \rightarrow (\mathcal{F}'_r)^{ab} = \mathcal{F}_r^{ab} \quad (3.5b)$$

$$r = 1, \dots, \lfloor \frac{\lambda-1}{2} \rfloor \quad (3.5c)$$

for any set of symmetric matrices $\Lambda_r^{ab} = \Lambda_r^{ba}$: If $\mathcal{L}_r^{ab}$ is a solution of (3.3) then $\mathcal{L}'_r^{ab}$ is also a solution.

Using the symmetry (2.14) again, we see that the gauge transformation (3.5) is also an invariance of the stress tensor

$$\hat{T} = \begin{cases} \mathcal{L}_0^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + \mathcal{L}_{\lambda/2}^{ab} : \hat{J}_a^{(\lambda/2)} \hat{J}_b^{(\lambda/2)} : + \sum_{r=1}^{\frac{\lambda}{2}-1} \mathcal{F}_r^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : & \text{for } \lambda \text{ even} \\ \mathcal{L}_0^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + \sum_{r=1}^{\frac{\lambda-1}{2}} \mathcal{F}_r^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : & \text{for } \lambda \text{ odd} \end{cases} \quad (3.6)$$

and similarly for the central charge. This tells us that all solutions in any given gauge orbit are physically equivalent, and we are entitled to choose a gauge.

3.3 The OVME

The most convenient gauge choice is

$$\mathcal{L}_r^{ab} = \mathcal{L}_{\lambda-r}^{ab}, \quad r = 1, \dots, \lfloor \frac{\lambda-1}{2} \rfloor \quad (3.7)$$

(choose $\Lambda_r^{ab} = (\mathcal{L}_{\lambda-r}^{ab} - \mathcal{L}_r^{ab})/2$) because this choice preserves the form of $\mathcal{L}_r^{ab}$ given for the twisted sectors of cyclic orbifolds in Ref. 9.

In this gauge we have

$$\mathcal{F}_r^{ab} = 2\mathcal{L}_r^{ab} \quad (3.8)$$

and the Virasoro condition (3.2) is summarized by the system

$$\hat{T} = \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (3.9a)$$

$$\mathcal{L}_r^{ab} = 2\mathcal{L}_r^{ac} \hat{G}_{cd} \mathcal{L}_r^{db} - \sum_{s=0}^{\lambda-1} \mathcal{L}_s^{cd} [\mathcal{L}_{r+s}^{ef} f_{ce}^a f_{df}^b + f_{ce}^f f_{df}^{(a} \mathcal{L}_r^{b)e}], \quad 0 \leq r \leq \lfloor \frac{\lambda}{2} \rfloor \quad (3.9b)$$

$$\mathcal{L}_r^{ab} = \mathcal{L}_{r \pm \lambda}^{ab} = \mathcal{L}_{\lambda \pm r}^{ab}, \quad \lfloor \frac{\lambda}{2} \rfloor < r \leq \lambda \quad (3.9c)$$

$$\hat{c} = 2\hat{G}_{ab} \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab}. \quad (3.9d)$$

In what follows, we will refer to (3.9b,c) as the *orbifold Virasoro master equation* (OVME).

As a check at $\lambda = 1$, we define

$$L^{ab} \equiv \mathcal{L}_0^{ab}, \quad J_a \equiv \hat{J}_a^{(0)}, \quad T \equiv \hat{T} \quad (3.10)$$

and the OVME reduces to the Virasoro master equation^{1,2} (VME):

$$T = L^{ab} : J_a J_b : \quad (3.11a)$$

$$L^{ab} = 2L^{ac} G_{cd} L^{db} - L^{cd} L^{ef} f_{ce}^a f_{df}^b - L^{cd} f_{ce}^f f_{df}^{(a} L^{b)e} \quad (3.11b)$$

$$G_{ab} = \oplus_I k_I \eta_{ab}^I, \quad c = 2G_{ab} L^{ab} \quad (3.11c)$$

as it should since $g_{\lambda=1}$ is affine Lie algebra.

The solutions of the OVME are in one-to-one correspondence with the conformal stress tensors $\hat{T}$ in (3.9a), and we find

$$L(m \geq 0)|0\rangle = \delta_{m,0} \hat{\Delta}_0 |0\rangle, \quad \hat{\Delta}_0 = \hat{G}_{ab} \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} \frac{r(\lambda-r)}{2\lambda^2} \quad (3.12)$$

where $\hat{\Delta}_0$ is the conformal weight of the ground state. This conformal weight is generically nonzero so that the generic construction of the OVME is like a twisted sector of an orbifold or a Ramond spin sector. The ground state conformal weight may occasionally vanish, e.g. the CFT's of the VME at $\lambda = 1$ have $\hat{\Delta}_0 = 0$.

Using the constraints (3.9c) to pull back the $\mathcal{L}$'s into the fundamental range^a,

$$\mathcal{L}_r^{ab}, \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \quad (3.13)$$

the OVME has the explicit form:

For even λ

$$\hat{T} = \mathcal{L}_0^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + \mathcal{L}_{\lambda/2}^{ab} : \hat{J}_a^{(\lambda/2)} \hat{J}_b^{(\lambda/2)} : + 2 \sum_{r=1}^{\frac{\lambda}{2}-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (3.14a)$$

$$\begin{aligned} \mathcal{L}_r^{ab} = & 2\mathcal{L}_r^{ac} \hat{G}_{cd} \mathcal{L}_r^{db} - (2 \sum_{s=0}^{\frac{\lambda}{2}-r-1} \mathcal{L}_s^{cd} \mathcal{L}_{r+s}^{ef} + \sum_{s=\frac{\lambda}{2}-r}^{\frac{\lambda}{2}} \mathcal{L}_s^{cd} \mathcal{L}_{\lambda-r-s}^{ef} + \sum_{s=0}^r \mathcal{L}_s^{cd} \mathcal{L}_{r-s}^{ef} - 2\mathcal{L}_0^{cd} \mathcal{L}_r^{ef}) f_{ce}^a f_{df}^b \\ & - (\mathcal{L}_0^{cd} + \mathcal{L}_{\lambda/2}^{cd} + 2 \sum_{s=1}^{\frac{\lambda}{2}-1} \mathcal{L}_s^{cd}) f_{ce}^f f_{df}^{(a} \mathcal{L}_r^{b)e}, \quad r = 0, \dots, \frac{\lambda}{2} \end{aligned} \quad (3.14b)$$

$$\hat{c} = 2\hat{G}_{ab} \mathcal{L}_0^{ab} + 2\hat{G}_{ab} \mathcal{L}_{\lambda/2}^{ab} + 4\hat{G}_{ab} \sum_{r=1}^{\frac{\lambda}{2}-1} \mathcal{L}_r^{ab} \quad (3.14c)$$

$$\hat{\Delta}_0 = \hat{G}_{ab} \mathcal{L}_{\lambda/2}^{ab} + \hat{G}_{ab} \sum_{r=1}^{\frac{\lambda}{2}-1} \mathcal{L}_r^{ab} \frac{r(\lambda-r)}{\lambda^2}. \quad (3.14d)$$

For odd λ

$$\hat{T} = \mathcal{L}_0^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + 2 \sum_{r=1}^{\frac{\lambda-1}{2}} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (3.15a)$$

$$\begin{aligned} \mathcal{L}_r^{ab} = & 2\mathcal{L}_r^{ac} \hat{G}_{cd} \mathcal{L}_r^{db} - (2 \sum_{s=1}^{\frac{\lambda-1}{2}-r} \mathcal{L}_s^{cd} \mathcal{L}_{r+s}^{ef} + \sum_{s=\frac{\lambda-1}{2}+1-r}^{\frac{\lambda-1}{2}} \mathcal{L}_s^{cd} \mathcal{L}_{\lambda-r-s}^{ef} + \sum_{s=0}^r \mathcal{L}_s^{cd} \mathcal{L}_{r-s}^{ef}) f_{ce}^a f_{df}^b \\ & - (\mathcal{L}_0^{cd} + 2 \sum_{s=1}^{\frac{\lambda-1}{2}} \mathcal{L}_s^{cd}) f_{ce}^f f_{df}^{(a} \mathcal{L}_r^{b)e}, \quad r = 0, \dots, \frac{\lambda-1}{2} \end{aligned} \quad (3.15b)$$

^aFor $\lambda = 1$ and 2 the fundamental range in (3.13) is the original range in (3.1b).

$$\hat{c} = 2\hat{G}_{ab}\mathcal{L}_0^{ab} + 4\hat{G}_{ab}\sum_{r=1}^{\frac{\lambda-1}{2}}\mathcal{L}_r^{ab} \quad (3.15c)$$

$$\hat{\Delta}_0 = \hat{G}_{ab}\sum_{r=1}^{\frac{\lambda-1}{2}}\mathcal{L}_r^{ab}\frac{r(\lambda-r)}{\lambda^2}. \quad (3.15d)$$

In these relations, we have used the convention that a sum is zero when its lower limit is greater than its upper limit. The forms (3.14b) and (3.15b) of the OVME are useful for computational purposes, while the form (3.9b) is useful for studying general properties of the system.

4 Properties of the OVME

4.1 Counting

The OVME is a set of

$$n(g, \lambda) = (\lfloor \frac{\lambda}{2} \rfloor + 1) \frac{\dim g(\dim g + 1)}{2} \quad (4.1)$$

coupled quadratic equations for the same number of unknowns (3.13).

The number of physically inequivalent solutions to the OVME expected³ at each level $\hat{k}$ is therefore

$$N(g, \lambda) \approx 2^{n(g, \lambda) - \dim g} \quad (4.2)$$

where we have subtracted the degrees of freedom associated with the Lie g covariance³

$$\mathcal{L}_r'^{ab} = \mathcal{L}_r^{cd}(\omega^{-1})_c^a(\omega^{-1})_d^b, \quad \omega \in Aut(g) \quad (4.3)$$

of the OVME. As examples of (4.2), one finds

$$N(SU(2), \lambda = 1) = 8, \quad N(SU(3), \lambda = 1) \approx \frac{1}{4} \text{ billion} \quad (4.4a)$$

$$N(SU(2), \lambda = 2) = 512, \quad N(SU(3), \lambda = 2) \approx 18 \text{ quintillion} \quad (4.4b)$$

where $\lambda = 1$ is the VME and the growth with λ at fixed g is exponential.

It is clear that, as in the VME, most of the solutions of the OVME will be new and (because the OVME is a large set of coupled quadratic equations) the new constructions will have generically irrational conformal weights and central charges.

4.2 $\mathcal{L}_r^{ab}$ independent of r

For any λ , we consider the simple consistent ansatz

$$\mathcal{L}_r^{ab}(\hat{G}) = \mathcal{L}^{ab}(\hat{G}), \quad (4.5)$$

($\hat{G}$ is defined in (2.3)) which collects all inverse inertia tensors independent of r . Then the OVME simplifies to a rescaled VME

$$\mathcal{L}^{ab}(\hat{G}) = \lambda[2\mathcal{L}^{ac}(\hat{G})G_{cd}\mathcal{L}^{db}(\hat{G}) - \mathcal{L}^{cd}(\hat{G})\mathcal{L}^{ef}(\hat{G})f_{ce}{}^a f_{df}{}^b - \mathcal{L}^{cd}(\hat{G})f_{ce}{}^f f_{df}{}^a \mathcal{L}^{be}(\hat{G})] \quad (4.6)$$

where G_{ab} is defined in (3.11c). Inspection of (4.6) gives

$$\hat{T} = \mathcal{L}^{ab}(\hat{G}) \sum_{r=0}^{\lambda-1} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (4.7a)$$

$$\mathcal{L}^{ab}(\hat{G}) = \frac{1}{\lambda} L^{ab}(G) \quad (4.7b)$$

$$\hat{c} = \lambda c = 2\lambda G_{ab} L^{ab}(G), \quad \hat{\Delta}_0 = \frac{\hat{c}}{24} \left(1 - \frac{1}{\lambda^2}\right) \quad (4.7c)$$

where $L^{ab}(G)$ is any solution of the VME (3.11b) on semisimple affine Lie g .

The constructions (4.7) were given in Ref. 9. They are the stress tensors of the twisted sectors of all the cyclic permutation orbifolds

$$\frac{A \times \dots \times A}{\mathbb{Z}_\lambda} \quad (4.8)$$

which can be constructed from λ copies of any affine-Virasoro construction (3.11a). In what follows, we will call (4.7) the set of *cyclic constructions*.^b

Because they operate in cyclic permutation orbifolds, the cyclic constructions form a Virasoro subalgebra of the orbifold Virasoro algebra⁹

$$\hat{T}^{(r)}(z)\hat{T}^{(s)}(w) = \frac{(\hat{c}/2)\delta_{r+s,0 \bmod \lambda}}{(z-w)^4} + \frac{2\hat{T}^{(r+s)}(w)}{(z-w)^2} + \frac{\partial_w \hat{T}^{(r+s)}(w)}{z-w} + O((z-w)^0) \quad (4.9a)$$

$$\hat{T}^{(r)}(z) = \frac{1}{\lambda} L^{ab}(G) \sum_{s=0}^{\lambda-1} : \hat{J}_a^{(s)}(z) \hat{J}_b^{(r-s)}(z) : \quad (4.9b)$$

^bThe stress tensors of the twisted sectors of the S_n permutation orbifolds can also be constructed^{15,9,16,17} as sums of commuting copies of the cyclic constructions (4.7). This is because every element of S_n can be expressed as a product of disjoint cyclic permutations.

where the zero-twist component $\hat{T}^{(0)}$ of the extended algebra is the cyclic construction $\hat{T}$ in (4.7).

Among the cyclic constructions, we note in particular the *orbifold affine-Sugawara construction* $\hat{T}_{g_\lambda}$,

$$\hat{T}_{g_\lambda} = \sum_{r=0}^{\lambda-1} (\mathcal{L}_r^{ab})_{g_\lambda} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (4.10a)$$

$$(\mathcal{L}_r^{ab})_{g_\lambda}(\hat{G}) = \frac{L_g^{ab}(G)}{\lambda} = \oplus_I \frac{\eta_I^{ab}}{2\hat{k}_I + \lambda Q_I} \quad (4.10b)$$

$$\hat{c}_{g_\lambda} = \lambda c_g = \lambda \sum_I \frac{2\hat{k}_I \dim g_I}{2\hat{k}_I + \lambda Q_I}, \quad \hat{\Delta}_0^{g_\lambda} = \frac{\hat{c}_{g_\lambda}}{24} \left(1 - \frac{1}{\lambda^2}\right) \quad (4.10c)$$

where Q_I is the quadratic Casimir of g_I and c_g is the central charge of the affine-Sugawara construction^{12,18–21} at levels $\{k_I\}$ on semisimple g . The form of these constructions^{13,14,9} on simple g is:

$$(\mathcal{L}_r^{ab})_{g_\lambda}(\hat{k}) = \frac{L_g^{ab}(k)}{\lambda} = \frac{\eta^{ab}}{2\hat{k} + \lambda Q_g} \quad (4.11a)$$

$$\hat{c}_{g_\lambda} = \frac{\lambda \hat{x} \dim g}{\hat{x} + \lambda \tilde{h}_g} = \lambda c_g(x), \quad c_g(x) = \frac{x \dim g}{x + \tilde{h}_g} \quad (4.11b)$$

$$\hat{\Delta}_0^{g_\lambda} = \frac{\hat{c}_{g_\lambda}}{24} \left(1 - \frac{1}{\lambda^2}\right). \quad (4.11c)$$

Here $\tilde{h}_g = Q_g/\psi^2$ is the dual Coxeter number of g , $x \in \mathbb{Z}^+$ is the invariant level in (2.4) and $c_g(x)$ is the central charge of the affine-Sugawara construction on simple g .

The orbifold affine-Sugawara constructions include the affine-Sugawara constructions T_g at $\lambda = 1$, and we shall see below that the orbifold affine-Sugawara constructions play the same fundamental role in the OVME that the affine-Sugawara constructions play in the VME.

4.3 Constructions on subalgebras

The OVME at order λ has solutions $\mathcal{L}(\lambda; h_\eta)$ given by

$$\mathcal{L}_r^{ab}(\lambda; h_\eta) = \begin{cases} \mathcal{L}_s^{AB}(h_\eta), & \text{if } \exists s \in [0, \eta - 1] \text{ s.t. } r = \frac{\lambda}{\eta} s \\ 0, & \text{otherwise} \end{cases} \quad (4.12a)$$

$$A, B = 1, \dots, \dim h, \quad r = 0, \dots, \lambda - 1, \quad \eta \in \mathbb{Z}^+, \quad \frac{\lambda}{\eta} \in \mathbb{Z}^+ \quad (4.12b)$$

$$\hat{c}(\lambda; h_\eta) = \hat{c}(h_\eta), \quad \hat{\Delta}_0(\lambda; h_\eta) = \hat{\Delta}_0(h_\eta) \quad (4.12c)$$

where $\mathcal{L}(h_\eta)$ is any solution to the OVME on h_η , $h \subset g$ (with central charge $\hat{c}(h_\eta)$ and ground state conformal weight $\hat{\Delta}_0(h_\eta)$). The order η solutions appear at order λ because $h_\eta \subset g_\lambda$ (see Section 2.2).

The subalgebra constructions (4.12) include in particular the orbifold affine-Sugawara construction on $h_\eta \subset g_\lambda$,

$$\hat{T}_{h_\eta}(\lambda; h_\eta) = \frac{\eta^{AB}}{2\hat{k} + \eta Q_h} \sum_{r=0}^{\eta-1} : \hat{J}_A^{(\frac{\lambda}{\eta} r)} \hat{J}_B^{(-\frac{\lambda}{\eta} r)} :, \quad A, B = 1, \dots, \dim h \quad (4.13a)$$

$$\hat{c}_{h_\eta} = \frac{r\eta \hat{x} \dim h}{r\hat{x} + \eta \hat{h}_h}, \quad \hat{\Delta}_0^{h_\eta} = \frac{\hat{c}_{h_\eta}}{24} \left(1 - \frac{1}{\eta^2}\right) \quad (4.13b)$$

for simple g and simple h where r is the index of embedding of $h \subset g$.

We also mention the case $h_\eta = g_1$ of (4.12), which collects all constructions on the integral affine subalgebra:

$$\mathcal{L}_r^{ab}(\lambda; g_1) = \delta_{r,0} L^{ab}(\hat{G}) \quad (4.14a)$$

$$\hat{T}(\lambda; g_1) = L^{ab}(\hat{G}) : \hat{J}_a^{(0)} \hat{J}_b^{(0)} :, \quad \hat{c}(\lambda; g_1) = 2L^{ab}(\hat{G}) \hat{G}_{ab}, \quad \hat{\Delta}_0(\lambda; g_1) = 0 \quad (4.14b)$$

where $L^{ab}(\hat{G})$ is any solution to the VME (3.11b) with G_{ab} replaced by $\hat{G}_{ab}$ in (2.3). These stress tensors are isomorphic to the stress tensors of the VME and the ground state conformal weights vanish, as expected, for all these constructions because $\mathcal{L}_0^{ab}$ does not contribute to $\hat{\Delta}_0$ in (3.12).

4.4 K-conjugation covariance

At order λ , the OVME exhibits *K-conjugation covariance* through the orbifold affine-Sugawara construction $\hat{T}_{g_\lambda}$, which includes the familiar K-conjugation covariance^{12,18,22,23,1} through the affine-Sugawara construction when $\lambda = 1$. This means (see App. A) that the K-conjugate inertia tensor $\tilde{\mathcal{L}}$

$$\tilde{\mathcal{L}}_r^{ab} = (\mathcal{L}_r^{ab})_{g_\lambda} - \mathcal{L}_r^{ab} \quad (4.15)$$

is a solution of the OVME when $\mathcal{L}$ is a solution, and

$$\tilde{\hat{T}} = \hat{T}_{g_\lambda} - \hat{T}, \quad \tilde{\hat{c}} = \hat{c}_{g_\lambda} - \hat{c}, \quad \tilde{\hat{\Delta}}_0 = \hat{\Delta}_0^{g_\lambda} - \hat{\Delta}_0, \quad (4.16a)$$

$$\hat{T}(z)\tilde{\hat{T}}(w) = O((z-w)^0) \quad (4.16b)$$

where $\tilde{\hat{T}}$ is the corresponding K-conjugate stress tensor.

Drawing on experience with the VME, K-conjugation covariance tells us that the conformal field theory corresponding to $\hat{T}$ can be considered as a gauge theory^{24,25,3,5-8} in which the cyclic permutation orbifold of λ copies of WZW is gauged by the K-conjugate partner $\tilde{\hat{T}}$ of $\hat{T}$.

The OVME has another covariance, which we call $Aut(\mathbb{Z}_\lambda)$ covariance, that has no analog in the VME: For any automorphism of $\mathbb{Z}_\lambda$, the inertia tensor $\mathcal{L}'$

$$\mathcal{L}'^{ab}_r = \mathcal{L}^{ab}_{\phi(r)}, \quad \hat{c}' = \hat{c}, \quad \phi \in Aut(\mathbb{Z}_\lambda) \quad (4.17)$$

is a solution when $\mathcal{L}$ is a solution. $Aut(\mathbb{Z}_\lambda)$ covariance relates conformal constructions with the same central charge but generically different conformal weights.

4.5 Coset constructions

At order λ , consider the orbifold affine-Sugawara construction $\hat{T}_{h_\eta} \equiv \hat{T}_{h_\eta}(\lambda; h_\eta)$ on $h_\eta \subset g_\lambda$ where g and h are semisimple. (For simple g and h , this construction is given in (4.13)). The K-conjugate partner of $\hat{T}_{h_\eta}$ is the general g_λ/h_η coset construction,

$$\hat{T}_{g_\lambda/h_\eta} = \tilde{\hat{T}} = \hat{T}_{g_\lambda} - \hat{T}_{h_\eta}, \quad \hat{c}_{g_\lambda/h_\eta} = \tilde{\hat{c}} = \hat{c}_{g_\lambda} - \hat{c}_{h_\eta} \quad (4.18)$$

which includes the ordinary coset constructions^{12,18,22} at $\lambda = 1$. The special case of (4.18) given by Kač and Wakimoto^{13,14} is $g_\lambda/g_{\eta=1}$, that is $\eta = 1$ and $h = g$ with

$$\hat{c}_{g_\lambda/g_{\eta=1}} = \hat{c}_{g_\lambda/g_1} = \lambda c_g(x) - c_g(\lambda x). \quad (4.19)$$

The extension to $g_\lambda/h_{\eta=1}$ was given in Ref. 9.

It was conjectured in Ref. 9 that the coset construction $g_\lambda/h_{\eta=1}$ corresponds to the twisted sectors of a different kind of orbifold

$$\frac{\left(\frac{g \times \dots \times g}{h_D(\lambda)} \right)^{\lambda \text{ times}}}{\mathbb{Z}_\lambda} \quad (4.20)$$

where $h_D(\lambda)$ is the diagonal subalgebra of $h \times \dots \times h$.

4.6 Nests

Repeated K-conjugation on nested orbifold affine subalgebras $g_\lambda \supset h_{\eta_1}^{(1)} \supset \dots \supset h_{\eta_n}^{(n)}$ gives the *orbifold affine-Sugawara nests*, which generalize the affine-Sugawara nests^{26,27,3} of the VME on $g \supset h^{(1)} \supset \dots \supset h^{(n)}$. The orbifold affine-Sugawara constructions and the coset constructions are the lowest nests, and the first non-trivial nests are

$$\hat{T}_{g_\lambda/h_\eta/h'_{\eta'}} = \hat{T}_{g_\lambda} - (\hat{T}_{h_\eta} - \hat{T}_{h'_{\eta'}}) \quad (4.21a)$$

$$\hat{c}_{g_\lambda/h_\eta/h'_{\eta'}} = \hat{c}_{g_\lambda} - (\hat{c}_{h_\eta} - \hat{c}_{h'_{\eta'}}). \quad (4.21b)$$

More generally one has the *orbifold affine-Virasoro nests*, including

$$\hat{T}_{g_\lambda/h_\eta/h'_{\eta'}}^\# = \hat{T}_{g_\lambda} - (\hat{T}_{h_\eta} - \hat{T}_{h'_{\eta'}}^\#) \quad (4.22a)$$

$$\hat{c}_{g_\lambda/h_\eta/h'_{\eta'}}^\# = \hat{c}_{g_\lambda} - (\hat{c}_{h_\eta} - \hat{c}_{h'_{\eta'}}^\#) \quad (4.22b)$$

where $\hat{T}_{h'_{\eta'}}^\#$ is an arbitrary construction on the subalgebra $h'_{\eta'}$. These nests generalize the affine-Virasoro nests^{27,3} familiar at $\lambda = 1$.

In what follows, we refer to the solutions discussed above (the cyclic constructions, the subalgebra constructions on $h_\eta \subset g_\lambda$ and the orbifold affine-Virasoro nests) as “known” solutions of the OVME and all other solutions will be called “new”.

5 The Lie g -Invariant Constructions

Many consistent ansätze^{27,3} and subansätze can be found for the OVME, as for the VME. In this section we concentrate on the Lie g -invariant constructions on simple g , whose abundance is a surprising feature of the OVME. The generalization of the “graph theory ansatz” on $SO(n)$, familiar²⁸ at $\lambda = 1$, is also given in App. B.

5.1 Group-invariant ansatz

We consider the group-invariant ansatz

$$\psi^2 \mathcal{L}_r^{ab} = l_r \eta^{ab}, \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \quad (5.1)$$

(η^{ab} is the inverse Killing metric of g) which collects all *Lie g -invariant constructions* in the OVME on simple g . We know that this ansatz includes at least the trivial construction

$\mathcal{L} = 0$, the orbifold affine-Sugawara construction $\hat{T}_{g_\lambda}$, the particular subalgebra constructions $\hat{T}_{g_\eta}$, the coset constructions g_λ/g_η and all orbifold affine-Sugawara nests of the form $g_\lambda \supset g_\eta \supset \dots \supset g_{\eta'}$. Among these only two constructions survive at $\lambda = 1$, namely the trivial construction and the affine-Sugawara construction on g . We shall see that the number of new Lie g -invariant constructions increases rapidly with λ .

Substitution of (5.1) into (3.9b) gives the reduced OVME of the ansatz:

$$\psi^2 \hat{T} = \sum_{r=0}^{\lambda-1} l_r \eta^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (5.2a)$$

$$l_r = \hat{x} l_r^2 + \tilde{h}_g \sum_{s=0}^{\lambda-1} l_s (2l_r - l_{r+s}), \quad 0 \leq r \leq \lfloor \frac{\lambda}{2} \rfloor \quad (5.2b)$$

$$l_r = l_{r \pm \lambda} = l_{\lambda \pm r}, \quad \lfloor \frac{\lambda}{2} \rfloor < r \leq \lambda \quad (5.2c)$$

$$\hat{c} = \hat{x} \dim g \left(\sum_{r=0}^{\lambda-1} l_r \right), \quad \hat{\Delta}_0 = \hat{x} \dim g \sum_{r=0}^{\lambda-1} l_r \frac{r(\lambda-r)}{4\lambda^2} \quad (5.2d)$$

where ψ is the highest root of g and $\hat{x} = \lambda x$, $x \in \mathbb{Z}^+$. For computational purposes we give also the forms of the reduced OVME in the fundamental range:

For even λ

$$\psi^2 \hat{T} = l_0 \eta^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + l_{\lambda/2} \eta^{ab} : \hat{J}_a^{(\lambda/2)} \hat{J}_b^{(-\lambda/2)} : + 2 \sum_{r=1}^{\frac{\lambda}{2}-1} l_r \eta^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (5.3a)$$

$$\begin{aligned} l_r = & \hat{x} l_r^2 - \tilde{h}_g \left(2 \sum_{s=0}^{\frac{\lambda}{2}-r-1} l_s l_{r+s} + \sum_{s=\frac{\lambda}{2}-r}^{\frac{\lambda}{2}} l_s l_{\lambda-r-s} + \sum_{s=0}^r l_s l_{r-s} - 2l_0 l_r \right) \\ & + 2\tilde{h}_g l_r (l_0 + l_{\lambda/2} + 2 \sum_{s=1}^{\frac{\lambda}{2}-1} l_s), \quad r = 0, \dots, \frac{\lambda}{2} \end{aligned} \quad (5.3b)$$

$$\hat{c} = \hat{x} \dim g (l_0 + l_{\lambda/2} + 2 \sum_{r=1}^{\frac{\lambda}{2}-1} l_r), \quad \hat{\Delta}_0 = \hat{x} \dim g \left(\frac{l_{\lambda/2}}{16} + \sum_{r=1}^{\frac{\lambda}{2}-1} l_r \frac{r(\lambda-r)}{2\lambda^2} \right). \quad (5.3c)$$

For odd λ

$$\psi^2 \hat{T} = l_0 \eta^{ab} : \hat{J}_a^{(0)} \hat{J}_b^{(0)} : + 2 \sum_{r=1}^{\frac{\lambda-1}{2}} l_r \eta^{ab} : \hat{J}_a^{(r)} \hat{J}_b^{(-r)} : \quad (5.4a)$$

$$\begin{aligned}
l_r = & \hat{x} l_r^2 - \tilde{h}_g \left(2 \sum_{s=1}^{\frac{\lambda-1}{2}-r} l_s l_{r+s} + \sum_{s=\frac{\lambda-1}{2}+1-r}^{\frac{\lambda-1}{2}} l_s l_{\lambda-r-s} + \sum_{s=0}^r l_s l_{r-s} \right) \\
& + 2\tilde{h}_g l_r (l_0 + 2 \sum_{s=1}^{\frac{\lambda-1}{2}} l_s), \quad r = 0, \dots, \frac{\lambda-1}{2}
\end{aligned} \tag{5.4b}$$

$$\hat{c} = \hat{x} \dim g(l_0 + 2 \sum_{r=1}^{\frac{\lambda-1}{2}} l_r), \quad \hat{\Delta}_0 = \hat{x} \dim g \left(\sum_{r=1}^{\frac{\lambda-1}{2}} l_r \frac{r(\lambda-r)}{2\lambda^2} \right). \tag{5.4c}$$

We see that the group-invariant ansatz (5.1) is a consistent ansatz, giving $\lfloor \frac{\lambda}{2} \rfloor + 1$ coupled quadratic equations and unknowns.

At each level $\hat{x}$, it follows that there are

$$N(\lambda) = 2^{\lfloor \frac{\lambda}{2} \rfloor + 1} \tag{5.5}$$

Lie g -invariant solutions at order λ . In fact, these solutions are organized into $N(\lambda)$ level families³, according to the high-level behavior^{28,3}

$$l_r = \frac{\theta_r}{\hat{x}} + O(\hat{x}^{-2}), \quad \theta_r \in \{0, 1\}, \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \tag{5.6}$$

of each level family. Only four of these level families are “known” at prime λ , and so there will be

$$N_{new}(\lambda) = 2^{\lfloor \frac{\lambda}{2} \rfloor + 1} - 4, \quad \lambda \text{ prime} \tag{5.7}$$

new Lie g -invariant level families at these orders.

The high-level expansion^{28,3} of the Lie g -invariant level families also shows that each level family is unitary for $\hat{x} = \lambda x$, $x \in \mathbb{Z}^+$, at least down to some finite radius of convergence $\hat{x}_0 = \lambda x_0$. Experience³ with the VME shows that x_0 is usually quite low, for example $x_0 = 1$ or 2 (see also Subsecs. 5.2 and 5.3).

We turn now to finding exact solutions of Eqs. (5.3b) and (5.4b). In solving these equations it was useful to note that only the $r = 0$ equations have l_0 dependence on their right hand sides, which effectively reduces the number of equations by one. Then we were able to factorize^{27,3} linear combinations of the $r \neq 0$ equations through $\lambda = 6$. We find no new solutions^c for $\lambda = 1, 2, 3$ and 4. The results for $\lambda = 5$ and 6 are reported below.

^cFor example the four solutions at $\lambda = 2$ are the trivial construction, the orbifold affine-Sugawara construction, the affine-Sugawara construction on the integral affine subalgebra and a Kač-Wakimoto coset.

5.2 Irrational conformal weights at $\lambda = 5$

At $\lambda = 5$ there are $2^{2+1} = 8$ level families, four of which are new:

$$l_0 = \frac{1}{2(\hat{x} + 5\tilde{h}_g)} + \frac{\theta}{2(\hat{x} + \tilde{h}_g)} \quad (5.8a)$$

$$l_1 = \frac{1}{2(\hat{x} + 5\tilde{h}_g)} (1 + \eta \sqrt{\frac{\hat{x} + 5\tilde{h}_g}{\hat{x} + \tilde{h}_g}}) \quad (5.8b)$$

$$l_2 = \frac{1}{2(\hat{x} + 5\tilde{h}_g)} (1 - \eta \sqrt{\frac{\hat{x} + 5\tilde{h}_g}{\hat{x} + \tilde{h}_g}}) \quad (5.8c)$$

$$\hat{c} = \frac{\hat{x}}{2} \dim g \left(\frac{5}{\hat{x} + 5\tilde{h}_g} + \frac{\theta}{\hat{x} + \tilde{h}_g} \right), \quad \eta, \theta = \pm 1. \quad (5.8d)$$

K-conjugation takes $(\theta, \eta) \rightarrow (-\theta, -\eta)$ and all these constructions are unitary ^d for $\hat{x} = 5x$, $x \in \mathbb{Z}^+$ because the $\mathcal{L}$'s are real³.

The central charges (5.8d) are rational, but the conformal weights of the ground states of these constructions are generically irrational

$$\hat{\Delta}_0 = \frac{\hat{x} \dim g}{10} \left(\frac{1}{\hat{x} + 5\tilde{h}_g} - \frac{\eta}{5\sqrt{(\hat{x} + 5\tilde{h}_g)(\hat{x} + \tilde{h}_g)}} \right) \quad (5.9)$$

and other conformal weights will be generically irrational because the $\mathcal{L}$'s are irrational. We will discuss this set of new constructions further in Sec. 6.

5.3 Unitary irrational central charge at $\lambda = 6$

At $\lambda = 6$ there are $2^{3+1} = 16$ level families, four of which are new:

$$l_0 = \frac{1}{2(\hat{x} + 6\tilde{h}_g)} \left(1 + \eta \frac{\tilde{h}_g(-\hat{x}^2 + 3\hat{x}\tilde{h}_g - 6\tilde{h}_g^2)}{(\hat{x} + \tilde{h}_g)\alpha} \right) + \frac{\theta}{2(\hat{x} + \tilde{h}_g)} \quad (5.10a)$$

$$l_1 = \frac{1}{2(\hat{x} + 6\tilde{h}_g)} \left(1 + \eta \frac{-\hat{x}^2 - 5\hat{x}\tilde{h}_g + 18\tilde{h}_g^2}{\alpha} \right) \quad (5.10b)$$

$$l_2 = \frac{1}{2(\hat{x} + 6\tilde{h}_g)} \left(1 + \eta \frac{\hat{x}^2 + 3\hat{x}\tilde{h}_g - 6\tilde{h}_g^2}{\alpha} \right) \quad (5.10c)$$

$$l_3 = \frac{1}{2(\hat{x} + 6\tilde{h}_g)} \left(1 + \eta \frac{\hat{x}^2 + \hat{x}\tilde{h}_g - 18\tilde{h}_g^2}{\alpha} \right) \quad (5.10d)$$

$$\alpha \equiv \sqrt{\hat{x}^4 + 2\hat{x}^3\tilde{h}_g - 19\hat{x}^2\tilde{h}_g^2 + 12\hat{x}\tilde{h}_g^3 + 36\tilde{h}_g^4}, \quad \eta, \theta = \pm 1. \quad (5.10e)$$

^dIt also follows that the high-level expansions of these level families are convergent down to and including level $x = 1$.

As in the previous case, K-conjugation takes $(\theta, \eta) \rightarrow (-\theta, -\eta)$.

Using the fact that the dual Coxeter number $\tilde{h}_g$ is positive for all compact Lie algebras, we have checked that α , and hence $\mathcal{L}_r^{ab}$, are real for all levels $\hat{x} = 6x$, $x \in \mathbb{Z}^+$ of all compact Lie algebras. It follows that these constructions are also unitary^e for these levels and algebras.

For these constructions, both the central charge and the ground state conformal weight are generically irrational:

$$\hat{c} = \frac{\hat{x}}{2} \dim g \left(\frac{6}{\hat{x} + 6\tilde{h}_g} + \eta \frac{\hat{x}^3 - 3\hat{x}^2\tilde{h}_g + 6\hat{x}\tilde{h}_g^2}{\alpha(\hat{x} + 6\tilde{h}_g)(\hat{x} + \tilde{h}_g)} + \frac{\theta}{\hat{x} + \tilde{h}_g} \right) \quad (5.11a)$$

$$\hat{\Delta}_0 = \frac{\hat{x} \dim g}{288(\hat{x} + 6\tilde{h}_g)} \left(35 + \eta \frac{15\hat{x}^2 + 7\hat{x}\tilde{h}_g - 78\tilde{h}_g^2}{\alpha} \right) \quad (5.11b)$$

due to the quantity α in (5.10e). For certain levels the central charge is rational (for example level $\hat{x} = 6x = 6$ of $SU(2)$ and $SU(3)$).

The lowest irrational central charges for each of the simple Lie algebras occur at $\theta = \eta = -1$ and are listed below:

	$\hat{x}$	$\hat{c}$
$SU(2)$	12	$\frac{9}{14}(5 - \frac{1}{\sqrt{2}}) \approx 2.7597$
$SO(3)$	12	$\frac{9}{14}(5 - \frac{1}{\sqrt{2}}) \approx 2.7597$
$SP(1)$	12	$\frac{9}{14}(5 - \frac{1}{\sqrt{2}}) \approx 2.7597$
E_6	6	$5 - \frac{19}{\sqrt{601}} \approx 4.2249$
E_7	6	$\frac{35}{8} - \frac{161}{8\sqrt{769}} \approx 3.6492$
E_8	6	$\frac{10}{3} - \frac{68}{3\sqrt{1471}} \approx 2.7423$
F_4	6	$\frac{26}{5} - \frac{26}{5\sqrt{46}} \approx 4.4330$
G_2	6	$\frac{21}{5} - \frac{21}{5\sqrt{41}} \approx 3.5440$

At fixed g , the central charges increase monotonically with $\hat{x}$, and, for the classical Lie algebras (e.g. $SU(n)$), the central charges increase monotonically with n .

The lowest unitary irrational central charge in this family of constructions is therefore

$$\hat{c}((E_8)_{\hat{x}=6}, \lambda = 6) = \frac{10}{3} - \frac{68}{3\sqrt{1471}} \approx 2.7423 \quad (5.12)$$

^eThe high-level expansions of these level families are also convergent through level $x = 1$.

and we note that level $\hat{x} = 6$ of E_8 corresponds, via the orbifold induction procedure⁹, to the historic level $x = 1$ of E_8 .

The value (5.12) is less than the lowest known unitary irrational central charge among the cyclic permutation orbifolds (4.8) at $\lambda = 6$,

$$\hat{c} = 6 \, c(SU(3)_5)_{D(1)}^\# = 6 \, (2)(1 - \frac{1}{\sqrt{61}}) \approx 10.4635 \quad (5.13)$$

where $c(SU(3)_5)_{D(1)}^\#$ is the lowest known³ unitary irrational central charge in the VME. The central charge in (5.12) is also less than $\hat{c} = 6 \times 1 = 6$, which is the lower bound on possible unitary irrational central charge of any order 6 cyclic permutation orbifold.

6 Discussion: The Lie h -invariant constructions

The solutions above for $\lambda \leq 6$ show that the Lie g -invariant constructions of the OVME are closely related to the Lie h -invariant constructions^{29,3}, with $h \subset g$, which have been studied in the VME. Following Ref. 29, the Lie h -invariant constructions of the OVME are those constructions on g_λ whose inertia tensors are Lie h -invariant

$$\delta \mathcal{L}_r^{ab} = \mathcal{L}_r^{c(a} f_{cd}^{b)} \psi^d = 0, \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \quad (6.1)$$

where ψ^a parameterizes the Lie group $H \subset G$ near the origin. Then we know from Ref. 29 that these constructions occur in Lie h -invariant quartets, octets, etc., and the currents of the Lie h -invariant stress tensors are either $(1, 0)$ operators (associated to a global h -symmetry) or $(0, 0)$ operators (associated to a local h -symmetry).

In the case of the OVME, Lie h symmetry guarantees at least a global or local h_1 symmetry, although h_η symmetry can also appear. For example, the general coset construction on g_λ/h_η lives in the Lie h -invariant quartet




Figure 1: The simplest Lie h -invariant quartet

where K_{g_λ} is the usual K-conjugation ($\hat{\tilde{T}} = \hat{T}_{g_\lambda} - \hat{T}$) through the orbifold affine-Sugawara construction on g_λ and K_{g_λ/h_η} is (a trivial example of) another K-conjugation through the coset construction itself ($\hat{\tilde{T}} = \hat{T}_{g_\lambda/h_\eta} - \hat{T}$ with $\hat{T} = 0$ or $\hat{T}_{g_\lambda/h_\eta}$).

The general Lie h -invariant quartet has the form

$$\begin{array}{ccc}
 \hat{T}(h_\eta \subset g_\lambda) & \xrightarrow{+\hat{T}_{h_\eta}} & \hat{T}(h_\eta \subset g_\lambda) + \hat{T}_{h_\eta} \\
 \uparrow K_{g_\lambda} & \swarrow K_{g_\lambda/h_\eta} & \uparrow K_{g_\lambda} \\
 \hat{T}_{g_\lambda} - \hat{T}(h_\eta \subset g_\lambda) & \xleftarrow{+\hat{T}_{h_\eta}} & \hat{T}_{g_\lambda/h_\eta} - \hat{T}(h_\eta \subset g_\lambda)
 \end{array}$$

Figure 2: The general Lie h -invariant quartet

where $\hat{T}(h_\eta \subset g_\lambda)$ is any locally Lie h -invariant construction on g_λ (so that $\hat{T}$ commutes with the currents of h_η). The K-conjugation K_{g_λ/h_η} through the coset construction relates the two locally-invariant constructions in the quartet, the other two constructions being globally invariant. For the Lie g -invariant ansatz, one should read $h_\eta \rightarrow g_\eta$.

We have checked that the two new sets of constructions at $\lambda = 5$ and 6 are Lie g -invariant quartets with the form

$$\begin{array}{ccc}
 \eta = +1, \theta = -1 & \xrightarrow{+\hat{T}_{g_1}} & \eta = +1, \theta = +1 \\
 \uparrow K_{g_\lambda} & \swarrow K_{g_\lambda/g_1} & \uparrow K_{g_\lambda} \\
 \eta = -1, \theta = +1 & \xleftarrow{+\hat{T}_{g_1}} & \eta = -1, \theta = -1
 \end{array}$$

Figure 3: New Lie g -invariant quartets at $\lambda = 5, 6$

so that K_{g_λ/g_1} is conjugation through the Kač-Wakimoto coset construction in both these cases. The constructions with $\theta = -1$ have a local g -invariance while the constructions with $\theta = 1$ have only a global g -invariance.

To check these conclusions for all the Lie g -invariant constructions, we have evaluated

the $\hat{T}\hat{J}^{(0)}$ OPE in this case to find

$$\hat{T}(z)\hat{J}_a^{(0)}(w) = M(\mathcal{L})_a{}^b \left(\frac{1}{(z-w)^2} + \frac{\partial_w}{z-w} \right) \hat{J}_b^{(0)} + O((z-w)^0) \quad (6.2a)$$

$$M(\mathcal{L})_a{}^b = \Delta \delta_a{}^b, \quad \Delta \equiv \hat{x}l_0 + \tilde{h}_g \sum_{r=0}^{\lambda-1} l_r \quad (6.2b)$$

where the matrix $M(\mathcal{L})_a{}^b$ is defined for all $\mathcal{L}$ in App. A. We have also checked that

$$\Delta^2 = \Delta \quad (6.3)$$

using the reduced OVME (5.2b) of the Lie g -invariant ansatz. The result (6.3) verifies that the currents $\hat{J}_a^{(0)}$ of the integral affine subalgebra $g_{\eta=1}$ are either $(1,0)$ or $(0,0)$ operators in all Lie g -invariant constructions, as they should be. The specific identifications in Fig. 3 follow easily from the central charges in (5.8d) and (5.11a).

Although all the Lie g -invariant constructions have a local or global symmetry associated to $g_{\eta=1}$, it is also possible to have larger symmetries associated to $h_\eta = g_\eta$, $\eta \geq 2$. Indeed, there is a quartet at $\lambda = 4$ associated to $g_{\eta=2}$. One remaining question is the nature of the extra (perhaps discrete) symmetry of $\mathcal{L}_r^{ab}$ which dictates the larger symmetry.

In the case of the $\lambda = 5$ quartet, one sees further structure because there are only two independent central charges in (5.8d). In fact, we may write these central charges as

$$\hat{c} = \frac{1}{2}(\hat{c}_{g_{\lambda=5}} + \theta \hat{c}_{g_{\eta=1}}) = \begin{cases} \frac{1}{2}(\hat{c}_{g_5} + \hat{c}_{g_1}) \\ \frac{1}{2}\hat{c}_{g_5/g_1} \end{cases} \quad (6.4)$$

which shows that the two g_1 -local theories ($\theta = -1$) have the same central charge, and moreover that this central charge is exactly half of the central charge of the Kač-Wakimoto coset.

Local Lie h -invariant constructions with $c = \frac{1}{2}c_{g/h}$, called the self $K_{g/h}$ -conjugate constructions^{29,3}, are known from the VME, where $\tilde{\hat{T}}$ and $\hat{T}$ in $\tilde{\hat{T}} = \hat{T}_{g_\lambda} - \hat{T}$ are automorphically equivalent inertia tensors (related by a transformation in $Aut(h)$). The mechanism here, although similar, is not the same: In the present case, conjugation by K_{g_5/g_1} ($\eta \rightarrow -\eta$) is an example of the $Aut(\mathbb{Z}_\lambda)$ covariance of the OVME (see Subsec. 4.4), so that the two constructions with $\hat{c} = \frac{1}{2}\hat{c}_{g_5/g_1}$ have generically different conformal weights.

The central charges in (5.8d) (or (6.4)) can also be written as

$$\hat{c} = 5\left(\frac{c_g(x)}{2}\right) + \theta\left(\frac{c_g(5x)}{2}\right), \quad c_g(x) = \frac{x \dim g}{x + \tilde{h}_g}, \quad x \in \mathbb{Z}^+ \quad (6.5)$$

where $c_g(x)$ is the central charge of the affine-Sugawara construction on g . Since conventional orbifoldization does not change the central charge of a theory, the form (6.5) suggests that the new constructions at $\lambda = 5$ might be twisted sectors of orbifolds which start with copies of a conformal field theory whose central charge is $c = c_g/2$. Conformal field theories with $c = c_g/2$ are in fact known (the self K-conjugate constructions^{28,3} at $\lambda = 1$), but these occur only for $\dim g = \text{even}$, leaving us without a conventional orbifold interpretation for the new constructions at $\lambda = 5$. For the new constructions at $\lambda = 6$, we have no suggestion at present for a conventional orbifold interpretation.

7 Extensions

In this section we consider the “Feigin-Fuchs” and the inner-automorphic deformations of the general Virasoro construction on orbifold affine algebra. For $\lambda \geq 3$, further deformations by the full antisymmetric part of the current bilinears $:\hat{J}_{[a}\hat{J}_{b]}:$ may also be possible.

7.1 $\hat{c}$ -changing deformations

The “Feigin-Fuchs” extension is

$$\hat{T}(z) = \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)}(z) \hat{J}_b^{(-r)}(z) : + D^a \partial \hat{J}_a^{(0)}(z) \quad (7.1a)$$

$$\mathcal{L}_r^{ab} = 2\mathcal{L}_r^{ac} \hat{G}_{cd} \mathcal{L}_r^{db} - \sum_{s=0}^{\lambda-1} \mathcal{L}_s^{cd} [\mathcal{L}_{r+s}^{ef} f_{ce}^a f_{df}^b + f_{ce}^f f_{df}^a (\mathcal{L}_r^{be})^c] + i f_{cd}^a (\mathcal{L}_r^{bc})^c D^d \quad (7.1b)$$

$$0 \leq r \leq \lfloor \frac{\lambda}{2} \rfloor \quad (7.1c)$$

$$\mathcal{L}_r^{ab} = \mathcal{L}_{r \pm \lambda}^{ab} = \mathcal{L}_{\lambda \pm r}^{ab}, \quad \lfloor \frac{\lambda}{2} \rfloor < r \leq \lambda \quad (7.1d)$$

$$D^a M(\mathcal{L})_a^b = D^b, \quad \hat{c} = 2\hat{G}_{ab} \left(\sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} - 6D^a D^b \right) \quad (7.1e)$$

where the matrix $M(\mathcal{L})_a^b$ is given in (A.7b). The ground state conformal weight $\hat{\Delta}_0$ is still given by (3.12), and these generalized $\hat{c}$ -changing deformations include the familiar c -changing deformations^{30,1} at $\lambda = 1$.

7.2 $\hat{c}$ -fixed deformations

Similarly, the generalized $\hat{c}$ -fixed deformations have the form

$$\hat{T}(z) = \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} : \hat{J}_a^{(r)}(z) \hat{J}_b^{(-r)}(z) : + d^a \frac{\hat{J}_a^{(0)}(z)}{z} + \frac{1}{2z^2} \hat{G}_{ab} d^a d^b \quad (7.2a)$$

$$d^a M(\mathcal{L})_a{}^b = d^b \quad (7.2b)$$

$$\hat{c} = 2\hat{G}_{ab} \sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab}, \quad \hat{\Delta}_0 = \frac{\hat{G}_{ab}}{2} \left(\sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab} \frac{r(\lambda-r)}{\lambda^2} + d^a d^b \right) \quad (7.2c)$$

and the OVME (3.9) holds for $\mathcal{L}_r^{ab}$. These deformations describe inner automorphic twists or spectral flow in the twisted sectors (see also Subsec. 7.4), a phenomenon which is familiar^{12,30,1} at $\lambda = 1$.

7.3 Deformation of the Lie g -invariant constructions

The Lie g -invariant constructions of Secs. 5 and 6 allow a large class of both the $\hat{c}$ -changing and $\hat{c}$ -fixed deformations. In this case we find that

$$f_{cd}{}^{(a} \mathcal{L}_r^{b)c} = 0 \quad (7.3a)$$

$$M(\mathcal{L})_a{}^b = \Delta \delta_a{}^b \rightarrow D^a(\Delta - 1) = 0 \quad (7.3b)$$

$$\Delta^2 = \Delta \rightarrow \Delta \in \{0, 1\} \quad (7.3c)$$

where Δ is given in (6.2b). The result (7.3a) tells us that the extra term in the generalized OVME (7.1b) vanishes, so that the reduced OVME (5.2b) is maintained for $\hat{c}$ -fixed or $\hat{c}$ -changing deformations of any Lie g -invariant construction.

We have recalled in (7.3c) that all currents $\hat{J}_a^{(0)}$ of any Lie g -invariant construction are either (1,0) or (0,0) operators. Then (7.3b) tells us that one may deform any Lie g -invariant construction by arbitrary D_a or d_a for any $\hat{J}_a^{(0)}$ which is a (1,0) current (global g symmetry) of the construction. In particular the orbifold affine-Sugawara construction may be deformed by arbitrary D or d , a fact which is familiar³⁰ for the affine-Sugawara constructions at $\lambda = 1$.

7.4 The doubly-twisted affine algebra

The $\hat{c}$ -fixed deformation of the orbifold affine-Sugawara construction for simple g

$$\hat{T}_{g_\lambda}(z; d) = \hat{T}_{g_\lambda}(z) + d^a \frac{\hat{J}_a^{(0)}(z)}{z} + \frac{1}{2z^2} \hat{k} \eta_{ab} d^a d^b, \quad \forall d \quad (7.4a)$$

$$\hat{c}_{g_\lambda}(d) = \lambda c_{g_\lambda}(x), \quad \hat{\Delta}_0^{g_\lambda}(d) = \frac{\hat{c}_{g_\lambda}}{24} \left(1 - \frac{1}{\lambda^2}\right) + \frac{1}{2} \hat{k} \eta_{ab} d^a d^b \quad (7.4b)$$

(see Eqs. (4.10) and (4.11)) describes spectral flow in the twisted sectors of the WZW permutation orbifolds

$$\frac{g(d) \overset{\lambda \text{ times}}{\times} \dots \times g(d)}{\mathbb{Z}_\lambda} \quad (7.5)$$

where each copy of g has been twisted by the same arbitrary vector d , using the known $\lambda = 1$ form³⁰ of (7.4).

Following Ref. 30, the spectral flow (7.4) is equivalent to introducing inner-automorphically twisted orbifold currents. In the Cartan-Weyl basis, these currents satisfy the “doubly-twisted” affine algebra

$$[\hat{H}_A^{(r)}(m + \frac{r}{\lambda}; d), \hat{H}_B^{(s)}(n + \frac{s}{\lambda}; d)] = \hat{k}(m + \frac{r}{\lambda}) \eta_{AB} \delta_{m+n+\frac{r+s}{\lambda}, 0} \quad (7.6a)$$

$$[\hat{H}_A^{(r)}(m + \frac{r}{\lambda}; d), \hat{E}_\alpha^{(s)}(n + \frac{s - d \cdot \alpha}{\lambda}; d)] = \alpha_A \hat{E}_\alpha^{(r+s)}(m + n + \frac{r + s - d \cdot \alpha}{\lambda}; d) \quad (7.6b)$$

$$[\hat{E}_\alpha^{(r)}(m + \frac{r - d \cdot \alpha}{\lambda}; d), \hat{E}_\beta^{(s)}(n + \frac{s - d \cdot \beta}{\lambda}; d)] \quad (7.6c)$$

$$= \begin{cases} N(\alpha, \beta) \hat{E}_{\alpha+\beta}^{(r+s)}(m + n + \frac{r+s-d \cdot (\alpha+\beta)}{\lambda}; d) & \text{if } (\alpha + \beta) \in \Delta \\ \alpha \cdot \hat{H}^{(r+s)}(m + n + \frac{r+s}{\lambda}; d) + \hat{k}(m + \frac{r-d \cdot \alpha}{\lambda}) \delta_{m+n+\frac{r+s}{\lambda}, 0} & \text{if } (\alpha + \beta) = 0 \\ 0 & \text{otherwise} \end{cases}$$

$$[\hat{H}_A^{(r)}(m + \frac{r}{\lambda}; d) - \delta_{m+\frac{r}{\lambda}, 0} k d_A] |0\rangle = \hat{E}_\alpha^{(r)}(m + \frac{r - d \cdot \alpha}{\lambda}; d) |0\rangle = 0 \quad \text{when } m + \frac{r}{\lambda} \geq 0 \quad (7.6d)$$

$$A, B = 1, \dots, \text{rank } g, \quad \alpha, \beta \in \Delta \quad (7.6e)$$

$$m, n \in \mathbb{Z}, \quad r, s = 0, \dots, \lambda - 1, \quad \lambda \in \mathbb{Z}^+ \quad (7.6f)$$

for all d_A , where $\hat{H}_A$ and $\hat{E}_\alpha$ are the Cartan and root operators respectively. The relations (7.6) are understood with the periodicity conditions

$$\hat{H}_A^{(r \pm \lambda)}(m + \frac{r \pm \lambda}{\lambda}; d) = \hat{H}_A^{(r)}(m \pm 1 + \frac{r}{\lambda}; d) \quad (7.7a)$$

$$\hat{E}_\alpha^{(r \pm \lambda)}(m + \frac{r \pm \lambda - d \cdot \alpha}{\lambda}; d) = \hat{E}_\alpha^{(r)}(m \pm 1 + \frac{r - d \cdot \alpha}{\lambda}; d) \quad (7.7b)$$

in analogy to the conditions (2.2).

The doubly-twisted algebra (7.6) reduces to the orbifold affine algebra (2.1) when $d = 0$ and it reduces to inner-automorphically twisted^{31,30,32} affine Lie algebra when $\lambda = 1$. More generally, the doubly-twisted algebra exhibits an intricate interplay between the discrete orbifold twist (outer automorphism) and the continuous spectral flow (inner automorphism).

The doubly-twisted algebra can be obtained by the orbifold induction procedure⁹ from the spectral flow at $\lambda = 1$ given in Ref. 30. The relations are

$$\begin{aligned}\hat{H}_A^{(r)}(m + \frac{r}{\lambda}; d) &\equiv \hat{J}_A^{(r)}(m + \frac{r}{\lambda}) + kd_A \delta_{m+\frac{r}{\lambda},0} \\ &= J_A(\lambda m + r; d)_{BOOST} \\ &= J_A(\lambda m + r) + kd_A \delta_{\lambda m+r,0}\end{aligned}\tag{7.8a}$$

$$\begin{aligned}\hat{E}_\alpha^{(r)}(m + \frac{r - d \cdot \alpha}{\lambda}; d) &\equiv \hat{E}_\alpha^{(r)}(m + \frac{r}{\lambda}) \\ &= E_\alpha(\lambda m + r - d \cdot \alpha; d)_{BOOST} \\ &= E_\alpha(\lambda m + r)\end{aligned}\tag{7.8b}$$

where the currents $J_A(m; d)_{BOOST}$, $E_\alpha(m - d \cdot \alpha; d)_{BOOST}$ are the inner automorphically-twisted currents of Ref. 30 and $J_A(m)$, $E_\alpha(m)$ satisfy affine Lie algebra.

Acknowledgements

We thank L. Loveridge and C. Schweigert for helpful discussions. J. E and M. B. H. were supported in part by the Director, Office of Energy Research, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract DE-AC03-76F00098 and in part by the National Science Foundation under grant PHY95-14797.

Appendix A. Derivations

Many of the operators below have branch points at zero and infinity; however we have checked that all the standard machinery of Ref. 33 follows using contours that never encircle these points.

•Wick's theorem

Given the OPE of two operators A and B ,

$$A(z)B(w) = \sum_m \frac{\{AB\}_m(w)}{(z-w)^m}\tag{A.1}$$

the contraction of the two operators is defined as the singular terms of the OPE

$$\underbrace{A(z)B(w)} \equiv \sum_{m>0} \frac{\{AB\}_m(w)}{(z-w)^m}. \quad (\text{A.2})$$

For any three operators A, B, C we have

$$\underbrace{A(z) : BC : (w)} = \oint_w \frac{(dx)}{x-w} \underbrace{A(z)B(x)} \underbrace{C(w)} + B(x) \underbrace{A(z)C(w)}. \quad (\text{A.3})$$

•Symmetry of the bilinears

To prove the symmetry (2.14) of the current bilinears, we use the fact that the commutator of the normal-ordered bilinear $: \hat{J} \hat{J} :$ is related to the derivatives of the singular terms of the $\hat{J} \hat{J}$ OPE (2.5a):

$$\begin{aligned} : \hat{J}_{(a)}^{(r)} \hat{J}_{(b)}^{(s)} : (z) - : \hat{J}_{(a)}^{(s)} \hat{J}_{(b)}^{(r)} : (z) &= \sum_{m>0} \frac{(-1)^{m+1}}{m!} \partial^m \{ \hat{J}_{(a)}^{(r)} \hat{J}_{(b)}^{(s)} \}_m(z) \\ &= \partial (i f_{(ab)}^c \hat{J}_c^{(r+s)}(z)) + \frac{-1}{2} \partial^2 (\hat{G}_{ab} \delta_{r+s,0 \bmod \lambda}) \\ &= 0. \end{aligned} \quad (\text{A.4})$$

In this paper, $A_{(a}B_{b)} = A_a B_b + A_b B_a$ and $A_{[a}B_{b]} = A_a B_b - A_b B_a$.

• $\hat{J} : \hat{J} \hat{J} :$ OPE

Starting with the $\hat{J} \hat{J}$ OPE (2.5a) and using Wick's theorem, we can evaluate the $\hat{J} : \hat{J} \hat{J} :$ OPE:

$$\begin{aligned} \underbrace{\hat{J}_a^{(r)}(z) : \hat{J}_b^{(s)} \hat{J}_c^{(t)} : (w)} &= \oint_w \frac{(dx)}{x-w} \underbrace{\hat{J}_a^{(r)}(z) \hat{J}_b^{(s)}(x)} \underbrace{\hat{J}_c^{(t)}(w)} + \hat{J}_b^{(s)}(x) \underbrace{\hat{J}_a^{(r)}(z) \hat{J}_c^{(t)}(w)} \\ &= \oint_w \frac{(dx)}{x-w} \left(\frac{\hat{G}_{ab} \delta_{r+s,0 \bmod \lambda}}{(z-x)^2} + \frac{i f_{ab}^d \hat{J}_d^{(r+s)}(x)}{(z-x)} \right) \hat{J}_c^{(t)}(w) \\ &\quad + \hat{J}_b^{(s)}(x) \left(\frac{\hat{G}_{ac} \delta_{r+t,0 \bmod \lambda}}{(z-w)^2} + \frac{i f_{ac}^d \hat{J}_d^{(r+t)}(w)}{(z-w)} \right) \\ &= \frac{i \hat{G}_{dc} f_{ab}^d \delta_{r+s+t,0 \bmod \lambda}}{(z-w)^3} \\ &\quad + \frac{(\hat{G}_{ab} \delta_c^e \delta_{r+s,0 \bmod \lambda} + \hat{G}_{ac} \delta_b^e \delta_{r+t,0 \bmod \lambda} - f_{ab}^d f_{dc}^e) \hat{J}_e^{(r+s+t)}(w)}{(z-w)^2} \\ &\quad + \frac{i f_{ab}^e : \hat{J}_e^{(r+s)} \hat{J}_c^{(t)} : (w) + i f_{ac}^d : \hat{J}_b^{(s)} \hat{J}_d^{(r+t)} : (w)}{z-w} \end{aligned}$$

$$\begin{aligned}
&= \frac{i\hat{G}_{ec}f_{ab}{}^e\delta_{r+s+t,0 \bmod \lambda}}{(z-w)^3} + \frac{M_{bca}^{(s,t,r)}{}^e\hat{J}_e^{(r+s+t)}(w)}{(z-w)^2} \\
&\quad + \frac{if_{ab}{}^e : \hat{J}_e^{(r+s)}\hat{J}_c^{(t)} : (w) + if_{ac}{}^e : \hat{J}_b^{(s)}\hat{J}_e^{(r+t)} : (w)}{z-w}
\end{aligned} \tag{A.5a}$$

$$M_{bca}^{(s,t,r)}{}^d \equiv \hat{G}_{ab}\delta_c^d\delta_{r+s,0 \bmod \lambda} + \hat{G}_{ac}\delta_b^d\delta_{r+t,0 \bmod \lambda} - f_{ab}{}^ef_{ec}{}^d. \tag{A.5b}$$

• : $\hat{J}\hat{J} : \hat{J}$ OPE

The : $\hat{J}\hat{J} : \hat{J}$ OPE is obtained by analytic continuation of (A.5a)

$$\begin{aligned}
\underbrace{: \hat{J}_a^{(r)}\hat{J}_b^{(s)} : (z)} \hat{J}_c^{(t)}(w) &= \frac{-i\hat{G}_{ec}f_{ab}{}^e\delta_{r+s+t,0 \bmod \lambda}}{(z-w)^3} \\
&\quad + \frac{M_{abc}^{(r,s,t)}{}^e\hat{J}_e^{(r+s+t)}(w)}{(z-w)^2} + \frac{M_{abc}^{(r,s,t)}{}^e\partial_w\hat{J}_e^{(r+s+t)}(w)}{(z-w)} \\
&\quad - \left(\frac{if_{ca}{}^e : \hat{J}_e^{(r+t)}\hat{J}_b^{(s)} : (w) + if_{cb}{}^e : \hat{J}_a^{(r)}\hat{J}_e^{(s+t)} : (w)}{z-w} \right). \tag{A.6}
\end{aligned}$$

• $\hat{T}\hat{J}^{(0)}$ OPE

Taking $s = -r$ and $t = 0$ and multiplying (A.6) by $\sum_{r=0}^{\lambda-1} \mathcal{L}_r^{ab}$, we obtain

$$\underbrace{\hat{T}(z)\hat{J}_a^{(0)}(w)} = M(\mathcal{L})_a{}^b \left(\frac{1}{(z-w)^2} + \frac{\partial_w}{z-w} \right) \hat{J}_b^{(0)}(w) + \frac{\sum_{r=0}^{\lambda-1} N^{(r)}(\mathcal{L})_a{}^{bc} : \hat{J}_b^{(r)}\hat{J}_c^{(-r)} : (w)}{z-w} \tag{A.7a}$$

$$M(\mathcal{L})_a{}^b \equiv \mathcal{L}_0^{cd}M_{cd,a}{}^b - \sum_{r=1}^{\lambda-1} \mathcal{L}_r^{cd}f_{ac}{}^ff_{fd}{}^b \quad N^{(r)}(\mathcal{L})_a{}^{bc} \equiv \mathcal{L}_r^{de}N_{de,a}{}^{bc} \tag{A.7b}$$

$$M_{ab,c}{}^d \equiv \delta_{(a}{}^d\hat{G}_{b)c} + \frac{1}{2}f_{e(a}{}^df_{b)c}{}^e \quad N_{ab,c}{}^{de} \equiv \frac{i}{2}\delta_{(a}^{(d}f_{b)c}{}^{e)}. \tag{A.7c}$$

• : $\hat{J}J :: \hat{J}J :$ OPE

To evaluate the OPE of two current bilinears we apply Wick's theorem again

$$\begin{aligned}
&\underbrace{: \hat{J}_a^{(r)}\hat{J}_b^{(s)} : (z)} : \hat{J}_c^{(t)}\hat{J}_d^{(u)} : (w) \\
&= \oint_w \frac{(dx)}{x-w} : \hat{J}_a^{(r)}\hat{J}_b^{(s)} : (z) \underbrace{\hat{J}_c^{(t)}(x)\hat{J}_d^{(u)}(w)} + \hat{J}_c^{(t)}(x) : \hat{J}_a^{(r)}\hat{J}_b^{(s)} : (z) \underbrace{\hat{J}_d^{(u)}(w)}
\end{aligned}$$

$$\begin{aligned}
&= \oint_w \frac{(dx)}{x-w} \left[\frac{-i\hat{G}_{ec} f_{ab}^e \delta_{r+s+t,0} \text{ mod } \lambda}{(z-x)^3} \right. \\
&\quad + \frac{M_{abc}^{(r,s,t)}{}^e \hat{J}_e^{(r+s+t)}(x)}{(z-x)^2} + \frac{M_{abc}^{(r,s,t)}{}^e \partial_x \hat{J}_e^{(r+s+t)}(x)}{(z-x)} \\
&\quad \left. - \left(\frac{if_{ca}^e : \hat{J}_e^{(r+t)} \hat{J}_b^{(s)} : (x) + if_{cb}^e : \hat{J}_a^{(r)} \hat{J}_e^{(s+t)} : (x)}{z-x} \right) \right] \hat{J}_d^{(u)}(w) \\
&\quad + \hat{J}_c^{(t)}(x) \left[\frac{-i\hat{G}_{ed} f_{ab}^e \delta_{r+s+u,0} \text{ mod } \lambda}{(z-w)^3} \right. \\
&\quad + \frac{M_{abd}^{(r,s,u)}{}^e \hat{J}_e^{(r+s+u)}(w)}{(z-w)^2} + \frac{M_{abd}^{(r,s,u)}{}^e \partial_w \hat{J}_e^{(r+s+u)}(w)}{(z-w)} \\
&\quad \left. - \left(\frac{if_{da}^e : \hat{J}_e^{(r+u)} \hat{J}_b^{(s)} : (w) + if_{db}^e : \hat{J}_a^{(r)} \hat{J}_e^{(s+u)} : (w)}{z-w} \right) \right] \tag{A.8a}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\hat{G}_{ed} M_{abc}^{(r,s,t)}{}^e \delta_{r+s+t+u,0} \text{ mod } \lambda}{(z-w)^4} \\
&\quad + \frac{-i\hat{G}_{ec} f_{ab}^e \delta_{r+s+t,0} \text{ mod } \lambda \hat{J}_d^{(u)}(w) - i\hat{G}_{ed} f_{ab}^e \delta_{r+s+u,0} \text{ mod } \lambda \hat{J}_c^{(t)}(w)}{(z-w)^3} \\
&\quad + \frac{(if_{fd}^e M_{abc}^{(r,s,t)}{}^f - if_{ca}^e M_{ebd}^{(r+t,s,u)}{}^f - if_{cb}^e M_{aed}^{(r,s+t,u)}{}^f) \hat{J}_f^{(r+s+t+u)}(w)}{(z-w)^3} \\
&\quad + \{ M_{abc}^{(r,s,t)}{}^e : \hat{J}_e^{(r+s+t)} \hat{J}_d^{(u)} : (w) + M_{abd}^{(r,s,u)}{}^e : \hat{J}_c^{(t)} \hat{J}_e^{(r+s+u)} : (w) \\
&\quad - f_{ca}^e (f_{de}^f : \hat{J}_f^{(r+t+u)} \hat{J}_b^{(s)} : (w) + f_{db}^f : \hat{J}_e^{(r+t)} \hat{J}_f^{(s+u)} : (w)) \\
&\quad - f_{cb}^e (f_{da}^f : \hat{J}_f^{(r+u)} \hat{J}_e^{(s+t)} : (w) + f_{de}^f : \hat{J}_a^{(r)} \hat{J}_f^{(s+t+u)} : (w)) \} / (z-w)^2 \\
&\quad + O((z-w)^{-1}). \tag{A.8b}
\end{aligned}$$

•Matching to the Virasoro algebra

We turn now to study the general stress tensor $\hat{T}$ in (3.1). To evaluate the $\hat{T}\hat{T}$ OPE set $s = -r$ and $u = -t$ in (A.8b) and multiply by $\sum_{r=0}^{\lambda-1} \sum_{t=0}^{\lambda-1} \mathcal{L}_r^{ab} \mathcal{L}_t^{cd}$. Comparing this to the Virasoro algebra (3.2) we find that the third-order pole term is zero since the identities

$$\sum_{r,s=0}^{\lambda-1} \mathcal{L}_r^{ab} \mathcal{L}_s^{cd} M_{aed}^{(r,-r+s,-s)}{}^f f_{cb}^e = 0. \tag{A.9a}$$

$$\mathcal{L}_r^{ab} \mathcal{L}_s^{cd} f_{fd}^e M_{abc}^{(r,-r,s)}{}^f = \mathcal{L}_r^{ab} \mathcal{L}_s^{cd} f_{ca}^e M_{ebd}^{(r+s,-r,-s)}{}^f \tag{A.9b}$$

follow by $a \leftrightarrow b$, $c \leftrightarrow d$ and $ab \leftrightarrow cd$ symmetry of $\sum_{r,s=0}^{\lambda-1} \mathcal{L}_r^{ab} \mathcal{L}_s^{cd}$.

Using the symmetry (2.14) the restriction (3.3) is obtained by matching the second-order pole terms. The fourth-order pole term gives the quadratic form of the central charge

$$\hat{c} = 2 \sum_{r,s=0}^{\lambda-1} \mathcal{L}_r^{ab} \mathcal{L}_s^{cd} \hat{G}_{ed} M_{abc}^{(r,-r,s) \ e} \quad (\text{A.10})$$

which can be simplified to the linear form in (3.4) using (3.3).

•First-order pole

The first-order pole term in the Virasoro algebra (3.2) is guaranteed to be correct because the higher-order pole terms are correct. To see this explicitly, we write

$$\hat{T}(z)\hat{T}(w) = \frac{\hat{c}/2}{(z-w)^4} + \frac{2\hat{T}(w)}{(z-w)^2} + \frac{F(w)}{z-w} + O((z-w)^0) \quad (\text{A.11})$$

where $F(w)$ is unknown, and this may be relabeled as

$$\hat{T}(w)\hat{T}(z) = \frac{\hat{c}/2}{(w-z)^4} + \frac{2\hat{T}(z)}{(w-z)^2} + \frac{F(z)}{w-z} + O((z-w)^0) \quad . \quad (\text{A.12})$$

We can determine F by a Taylor expansion $z = w + (z-w)$ of (A.11) and its analytic continuation $\hat{T}(z)\hat{T}(w) = \hat{T}(w)\hat{T}(z)$:

$$\hat{T}(w)\hat{T}(z) = \frac{\hat{c}/2}{(z-w)^4} + \frac{2\hat{T}(z)}{(z-w)^2} + \frac{2\partial_z \hat{T}(z) - F(z)}{w-z} + O((z-w)^0) \quad . \quad (\text{A.13})$$

Comparing to (A.12) we see that

$$F(w) = \partial_w \hat{T}(w) \quad (\text{A.14})$$

which completes the verification of the Virasoro algebra (3.2).

•K-conjugation

In parallel with the VME, it is straightforward to check that the K-conjugate partner $\tilde{\mathcal{L}}_r^{ab} = (\mathcal{L}_r^{ab})_{g_\lambda} - \mathcal{L}_r^{ab}$ of any solution $\mathcal{L}_r^{ab}$ of the OVME is also a solution of the OVME (use the fact that both $(\mathcal{L}_r^{ab})_{g_\lambda}$ and $\mathcal{L}_r^{ab}$ are solutions).

To see that the stress tensors of K-conjugate constructions commute, we use (A.8b) to verify the OPE

$$\begin{aligned} \hat{T}_{g_\lambda}(z)\hat{T}(w) &= \frac{\hat{c}/2}{(z-w)^4} + \frac{2\hat{T}(w)}{(z-w)^2} + \frac{\partial_w \hat{T}(w)}{(z-w)} + O((z-w)^0) \\ &= \hat{T}(z)\hat{T}(w) + O((z-w)^0) \end{aligned} \quad (\text{A.15})$$

and this is easily rearranged into (4.16b).

Appendix B. Graph Theory Ansatz on $g_\lambda = SO(n)_\lambda$

The graph theory ansatz $SO(n)_{diag}$ on $SO(n)_{\lambda=1}$ is familiar^{28,3} from the VME. Here we find the generalization in the OVME to the ansatz $SO(n)_\lambda^{diag}$ on $SO(n)_\lambda$.

The inverse inertia tensor in this case has the form

$$\psi^2 \mathcal{L}_r^{ab} = \psi^2 \mathcal{L}_r^{ij,kl} = \eta^{ij,kl} \mathcal{L}_r^{ij} = \delta^{ik} \delta^{jl} \mathcal{L}_r^{ij}, \quad a = (i, j), \quad 1 \leq i < j \leq n \quad (\text{B.1})$$

in the vector index notation for the Cartesian basis of $SO(n)$. Substituting into the OVME on $g_\lambda = SO(n)_\lambda$, we find the reduced system

$$\psi^2 \hat{T} = \sum_{r=0}^{\lambda-1} \sum_{i < j} \mathcal{L}_r^{ij} : \hat{J}_{ij}^{(r)} \hat{J}_{ij}^{(-r)} : \quad (\text{B.2a})$$

$$\mathcal{L}_r^{ij} (1 - \hat{x} \mathcal{L}_r^{ij}) + \tau_n \sum_{s=0}^{\lambda-1} \sum_{l \neq i,j}^n [\mathcal{L}_s^{il} \mathcal{L}_{r+s}^{lj} - \mathcal{L}_r^{ij} (\mathcal{L}_s^{il} + \mathcal{L}_s^{jl})] = 0 \quad (\text{B.2b})$$

$$\mathcal{L}_r^{ij} = \mathcal{L}_{\lambda \pm r}^{ij} = \mathcal{L}_{r \pm \lambda}^{ij}, \quad \lfloor \frac{\lambda}{2} \rfloor < r \leq \lambda \quad (\text{B.2c})$$

$$\mathcal{L}_r^{ij} = \mathcal{L}_r^{ji}, \quad \mathcal{L}_r^{ii} \equiv 0, \quad \tau_n = \begin{cases} 1, & n \neq 3 \\ 2, & n = 3 \end{cases} \quad (\text{B.2d})$$

$$\hat{c} = \hat{x} \sum_{r=0}^{\lambda-1} \sum_{i < j} \mathcal{L}_r^{ij}, \quad \hat{\Delta}_0 = \hat{x} \sum_{r=0}^{\lambda-1} \sum_{i < j} \mathcal{L}_r^{ij} \frac{r(\lambda - r)}{4\lambda^2} \quad (\text{B.2e})$$

which further reduces to $SO(n)_{diag}$ when $\lambda = 1$. The conventions in (B.2d) follow Ref. 28.

The reduced OVME in (B.2) consists of $(\lfloor \frac{\lambda}{2} \rfloor + 1) \binom{n}{2}$ equations and unknowns, so the reduced system contains

$$N(SO(n)_{diag}, \lambda) = 2^{(\lfloor \frac{\lambda}{2} \rfloor + 1) \binom{n}{2}} \quad (\text{B.3})$$

level families for each value of n and λ . As in the case of the Lie g -invariant constructions, most of these level families will be new, with irrational central charges and (for $\lambda \geq 2$) generically irrational ground state conformal weights $\hat{\Delta}_0$.

These level families can be classified by the graphs G_n^λ of order n with $\lfloor \frac{\lambda}{2} \rfloor + 1$ colors, according to the high-level behavior^{28,3}:

$$\mathcal{L}_r^{ij}(G_n^\lambda) = \frac{\Theta_r^{ij}(G_n^\lambda)}{\hat{x}} + O(\hat{x}^{-2}), \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \quad (\text{B.4a})$$

$$\psi^2 \hat{T}(G_n^\lambda) = \sum_{r=0}^{\lambda-1} \sum_{i < j} \left(\frac{\Theta_r^{ij}(G_n^\lambda)}{\hat{x}} + O(\hat{x}^{-2}) \right) : \hat{J}_{ij}^{(r)} \hat{J}_{ij}^{(-r)} : \quad (\text{B.4b})$$

$$\hat{c}(G_n^\lambda) = \sum_{r=0}^{\lambda-1} \sum_{i < j} \Theta_r^{ij}(G_n^\lambda) + O(\hat{x}^{-1}) \quad (\text{B.4c})$$

$$\hat{\Delta}_0(G_n^\lambda) = \sum_{r=0}^{\lambda-1} \sum_{i < j} \Theta_r^{ij} \frac{r(\lambda - r)}{4\lambda^2} + O(\hat{x}^{-1}) \quad (\text{B.4d})$$

of each level family, where

$$\Theta_r^{ij}(G_n^\lambda) \in \{0, 1\}, \quad 1 \leq i < j \leq n, \quad r = 0, \dots, \lfloor \frac{\lambda}{2} \rfloor \quad (\text{B.5})$$

is the adjacency matrix of any colored graph G_n^λ . The points i and j of G_n^λ are connected by an edge of color r when $\Theta_r^{ij}(G_n^\lambda) = 1$, and the collection of edges between any i and j is called a bond.

The high-level expansion also shows that each level family is unitary for $\hat{x} = \lambda x$, $x \in \mathbb{Z}^+$, at least down to some finite radius of convergence $\hat{x}_0 = \lambda x_0$.

In the colored graph theory, the orbifold affine-Sugawara construction $SO(n)_\lambda$ lives on the complete colored graph with n points and $\lfloor \frac{\lambda}{2} \rfloor + 1$ colors in each bond. As an example, the graph of the orbifold affine-Sugawara construction $SO(3)_{\lambda=2}$ is shown (with colors 0 and 1) below




Figure 4: The orbifold affine-Sugawara construction $SO(3)_{\lambda=2}$

so that $\hat{c} = 6 + O(\hat{x}^{-1})$ in this case.

Many other properties of the level families in this ansatz can be seen in the colored graphs, in analogy with the application²⁸ of ordinary graph theory at $\lambda = 1$. As examples,

- $\text{Aut}(\text{SO}(n))$ -inequivalent level families are in one-to-one correspondence with the (point) unlabelled colored graphs.

- Since K-conjugation is through the orbifold affine-Sugawara constructions, K-conjugate level families live on complementary colored graphs.

- Self K-conjugate constructions^{28,3}, with half the orbifold affine-Sugawara central charge

$$\hat{c} = \frac{\hat{c}_{g_\lambda}}{2} \quad (\text{B.6})$$

live on self-complementary colored graphs.

References

- [1] M. B. Halpern and E. Kiritsis, *Mod. Phys. Lett.* **A4**, 1373 (1989); Erratum, *ibid.* **A4**, 1797 (1989).
- [2] A. Yu Morozov, A. M. Perelomov, A. A. Rosly, M. A. Shifman and A. V. Turbiner, *Int. J. Mod. Phys.* **A5**, 803 (1990).
- [3] M. B. Halpern, E. Kiritis, N. A. Obers and K. Clubok, “Irrational Conformal Field Theory,” *Physics Reports* **265** No. 1&2 (1996).
- [4] M. B. Halpern and N. A. Obers, *Int. J. Mod. Phys.* **A11**, 4837 (1996).
- [5] J. de Boer and M. B. Halpern, *Int. J. Mod. Phys.* **A12**, 1551 (1997).
- [6] J. de Boer and M. B. Halpern, “Unification of the General Non-Linear Sigma Model and the Virasoro Master Equation”. In the proceedings of the NATO Workshop “New Developments in Quantum Field Theory” Zakopane, Poland (1997).
- [7] J. de Boer and M. B. Halpern, *Theor. and Math. Phys.* **V.117**, n.2, 1286 (1998).
- [8] J. de Boer and M. B. Halpern, *Int. J. Mod. Phys.* **A13**, 4487 (1998).
- [9] L. Borisov, M. B. Halpern and C. Schweigert, *Int. J. Mod. Phys.* **A13**, 125 (1998).
- [10] V. G. Kač, *Funct. Anal. App.* **1**, 328 (1967).
- [11] R. V. Moody, *Bull. Am. Math. Soc.* **73**, 217 (1967).
- [12] K. Bardakçi and M. B. Halpern, *Phys. Rev.* **D3**, 2493 (1971).
- [13] V. G. Kač and M. Wakimoto, *Acta Appl. Math.* **21**, 3 (1990).
- [14] P. Bouwknegt, “A New Coset Construction and Applications,” q-alg/9610013.
- [15] R. Dijkgraaf, G. Moore, E. Verlinde and H. Verlinde, *Comm. Math. Phys.* **185**, 197 (1997).
- [16] P. Bantay, *Phys. Lett.* **B419**, 175 (1998).
- [17] C. Schweigert, “Algebraic Aspects of Orbifold Theories”. Unpublished talk given at the workshop “Beyond the Standard Model”, Bad Honnef (1997).
- [18] M. B. Halpern, *Phys. Rev.* **D4**, 2398 (1971).
- [19] R. Dashen and Y. Frishman, *Phys. Rev.* **D11**, 278 (1975).
- [20] V. G. Knizhnik and A. B. Zamolodchikov, *Nucl. Phys.* **B247**, 83 (1984).
- [21] G. Segal, unpublished.
- [22] P. Goddard, A. Kent and D. Olive, *Phys. Lett.* **B152**, 88 (1985).
- [23] E. Kiritsis, *Mod. Phys. Lett.* **A4**, 437 (1989).
- [24] M. B. Halpern and J. P. Yamron, *Nucl. Phys.* **B351**, 333 (1991).
- [25] J. de Boer, K. Clubok and M. B. Halpern, *Int. J. Mod. Phys.* **A9**, 2451 (1994).
- [26] E. Witten, “The Central Charge in Three Dimensions”, in Memorial Volume for V. Knizhnik, ed. L. Brink et. al., *World Scientific* (1990).

- [27] M. B. Halpern, E. Kiritsis, N. A. Obers, M. Porrati and J. P. Yamron, *Int. J. Mod. Phys. A* **5**, 2275 (1990).
- [28] M. B. Halpern and N. A. Obers, *Comm. Math. Phys.* **138**, 63 (1991).
- [29] M. B. Halpern, E. Kiritsis and N. A. Obers, *Int. J. Mod. Phys. A* **7**, [Suppl.1A] 339 (1992).
- [30] J. K. Freericks and M. B. Halpern, *Ann. of Phys.* **188**, 258 (1988); Erratum, *ibid.* **190**, 212 (1989).
- [31] P. Goddard and D. Olive, *Int. J. Mod. Phys. A* **1**, 303 (1986).
- [32] W. Lerche, C. Vafa and N. P. Warner, *Nucl. Phys.* **B324**, 427 (1989).
- [33] F. A. Bais, P. Bouwknegt, K. Schoutens and M. Surridge, *Nucl. Phys.* **B304**, 348 (1988).